

**STATEMENT CLAIMING SMALL ENTITY STATUS
(37 CFR 1.9(f) & 1.27(c)) -- SMALL BUSINESS CONCERN**

Docket Number (Optional)
LEX-0071-USA

Applicant, Patentee, or Identifier: Gregory Donoho et al.

Application or Patent No.: _____

Filed or Issued: October 18, 2000

Title: Novel Human Proteins and Polynucleotides Encoding the Same

I hereby state that I am

- ☐ the owner of the small business concern identified below;
☒ an official of the small business concern empowered to act on behalf of the concern identified below:

NAME OF SMALL BUSINESS CONCERN Lexicon Genetics Incorporated

ADDRESS OF SMALL BUSINESS CONCERN 4000 Research Forest Drive, The Woodlands, TX 77381

I hereby state that the above identified small business concern qualifies as a small business concern as defined in 13 CFR Part 121 for purposes of paying reduced fees to the United States Patent and Trademark Office. Questions related to size standards for a small business concern may be directed to: Small Business Administration, Size Standards Staff, 409 Third Street, SW, Washington, DC 20416.

I hereby state that rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the invention described in:

- ☒ the specification filed herewith with title as listed above.
☐ the application identified above.
☐ the patent identified above.

If the rights held by the above identified small business concern are not exclusive, each individual, concern, or organization having rights in the invention must file separate statements as to their status as small entities, and no rights to the invention are held by any person, other than the inventor, who would not qualify as an independent inventor under 37 CFR 1.9(c) if that person made the invention, or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d), or a nonprofit organization under 37 CFR 1.9(e).

Each person, concern, or organization having any rights in the invention is listed below:

- ☒ No such persons, concerns, or organizations exist.
☐ each such person, concern, or organization is listed below:

Separate statements are required from each named person, concern, or organization having rights to the invention stating their status as small entities. (37 CFR 1.27)

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))

NAME OF PERSON SIGNING Lance K. Ishimoto

TITLE OF PERSON IF OTHER THAN OWNER Vice President - Intellectual Property

ADDRESS OF PERSON SIGNING 4000 Research Forest Drive, The Woodlands, TX 77381

SIGNATURE _____

DATE October 18, 2000

Reg. No. 41,866

**NOVEL HUMAN PROTEINS AND
POLYNUCLEOTIDES ENCODING THE SAME**

The present invention claims the benefit of U.S. Provisional
5 Application Numbers 60/160,285 and 60/183,583 which were filed
October 19, 1999 and February 18, 2000 respectively and are herein
incorporated in their entirety.

1. INTRODUCTION

10 The present invention relates to the discovery,
identification, and characterization of novel human
polynucleotides encoding proteins sharing sequence similarity with
mammalian proteins having CUB domains. The invention encompasses
the described polynucleotides, host cell expression systems, the
15 encoded proteins, fusion proteins, polypeptides and peptides,
antibodies to the encoded proteins and peptides, and genetically
engineered animals that either lack or over express the disclosed
sequences, antagonists and agonists of the proteins, and other
compounds that modulate the expression or activity of the proteins
20 encoded by the disclosed polynucleotides that can be used for
diagnosis, drug screening, clinical trial monitoring, or the
treatment of physiological disorders, or diseases.

2. BACKGROUND OF THE INVENTION

25 The CUB domain is an extracellular domain (ECD) present in
variety of diverse proteins such as bone morphogenetic protein 1,
proteinases, spermadhesins, complement subcomponents, and neuronal
recognition molecules. Given the importance of these functions,
CUB proteins have been associated with, *inter alia*, regulating
30 development, modulating cellular processes, and preventing
infectious disease.

3. SUMMARY OF THE INVENTION

The present invention relates to the discovery, identification, and characterization of nucleotides that encode novel human proteins, and the corresponding amino acid sequences of these proteins. The novel human proteins (NHPs) described for the first time herein share structural similarity with animal CUB domain proteins, coagulation factors V and XIII, milk fat globule-EGF factor 8, transcriptional repressor AE-binding protein-1, and neuropilins 1 and 2 (which, like the presently described protein, contain both CUB and discoidin domains).

The novel human nucleic acid (cDNA) sequences described herein, encode proteins/open reading frames (ORFs) of 487, 586, and 539 amino acids in length (see SEQ ID NOS: 2, 4, and 6 respectively).

The invention also encompasses agonists and antagonists of the described NHPs, including small molecules, large molecules, mutant NHPs, or portions thereof that compete with native NHPs, NHP peptides, and antibodies, as well as nucleotide sequences that can be used to inhibit the expression of the described NHPs (e.g., antisense and ribozyme molecules, and gene or regulatory sequence replacement constructs) or to enhance the expression of the described NHPs (e.g., expression constructs that place the described gene under the control of a strong promoter system), and transgenic animals that express a NHP transgene, or "knock-outs" (which can be conditional) that do not express a functional NHP. Several knockout ES cell lines have been produced that contain a gene trap mutation in a murine ortholog/homolog of the disclosed NHPs.

Further, the present invention also relates to processes for identifying compounds that modulate, i.e., act as agonists or antagonists, of NHP expression and/or NHP activity that utilize purified preparations of the described NHPs and/or NHP products, or cells expressing the same. Such compounds can be used as

therapeutic agents for the treatment of any of a wide variety of symptoms associated with biological disorders or imbalances.

4. DESCRIPTION OF THE SEQUENCE LISTING AND FIGURES

5 The Sequence Listing provides the sequences of several NHP ORFs encoding the described NHP amino acid sequences. SEQ ID NO:7 describes a NHP ORF and flanking sequences.

5. DETAILED DESCRIPTION OF THE INVENTION

10 The NHPs described for the first time herein are novel proteins that are expressed in, *inter alia*, human cell lines, and human prostate, pituitary, fetal brain, brain, thymus, spleen, lymph node, trachea, kidney, fetal liver, thyroid, adrenal gland, salivary gland, stomach, small intestine, colon, muscle, heart, 15 mammary gland, adipose, skin, esophagus, bladder, cervix, rectum, and testis cells.

The described sequences were compiled from gene trapped cDNAs, genomic sequence, and clones isolated from human brain, adipose, testis, and placenta cDNA libraries (Edge Biosystems, Gaithersburg, MD, and Clontech, Palo Alto, CA). The present 20 invention encompasses the nucleotides presented in the Sequence Listing, host cells expressing such nucleotides, the expression products of such nucleotides, and: (a) nucleotides that encode mammalian homologs of the described genes, including the 25 specifically described NHPs, and NHP products; (b) nucleotides that encode one or more portions of a NHP that correspond to functional domains, and the polypeptide products specified by such nucleotide sequences, including but not limited to the novel regions of any active domain(s); (c) isolated nucleotides that 30 encode mutant versions, engineered or naturally occurring, of the described NHPs in which all or a part of at least one domain is deleted or altered, and the polypeptide products specified by such nucleotide sequences, including but not limited to soluble

proteins and peptides in which all or a portion of the signal sequence is deleted; (d) nucleotides that encode chimeric fusion proteins containing all or a portion of a coding region of a NHP, or one of its domains (e.g., a receptor or ligand binding domain, 5 accessory protein/self-association domain, etc.) fused to another peptide or polypeptide; or (e) therapeutic or diagnostic derivatives of the described polynucleotides such as oligonucleotides, antisense polynucleotides, ribozymes, dsRNA, or gene therapy constructs comprising a sequence first disclosed in 10 the Sequence Listing.

As discussed above, the present invention includes: (a) the human DNA sequences presented in the Sequence Listing (and vectors comprising the same) and additionally contemplates any nucleotide sequence encoding a contiguous NHP open reading frame (ORF), or a 15 contiguous exon splice junction first described in the Sequence Listing, that hybridizes to a complement of a DNA sequence presented in the Sequence Listing under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and 20 washing in 0.1xSSC/0.1% SDS at 68°C (Ausubel F.M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York, at p. 2.10.3) and encodes a functionally equivalent gene product. Additionally contemplated are any nucleotide sequences 25 that hybridize to the complement of the DNA sequence that encode and express an amino acid sequence presented in the Sequence Listing under moderately stringent conditions, e.g., washing in 0.2xSSC/0.1% SDS at 42°C (Ausubel et al., 1989, *supra*), yet still encode a functionally equivalent NHP product. Functional 30 equivalents of a NHP include naturally occurring NHPs present in other species and mutant NHPs whether naturally occurring or engineered (by site directed mutagenesis, gene shuffling, directed evolution as described in, for example, U.S. Patent No.

5,837,458). The invention also includes degenerate nucleic acid variants of the disclosed NHP polynucleotide sequences.

Additionally contemplated are polynucleotides encoding a NHP ORF, or its functional equivalent, encoded by a polynucleotide sequence that is about 99, 95, 90, or about 85 percent similar or identical to corresponding regions of the nucleotide sequences of the Sequence Listing (as measured by BLAST sequence comparison analysis using, for example, the GCG sequence analysis package using standard default settings).

The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore the complements of, the described NHP gene nucleotide sequences. Such hybridization conditions may be highly stringent or less highly stringent, as described above. In instances where the nucleic acid molecules are deoxyoligonucleotides ("DNA oligos"), such molecules are generally about 16 to about 100 bases long, or about 20 to about 80, or about 34 to about 45 bases long, or any variation or combination of sizes represented therein that incorporate a contiguous region of sequence first disclosed in the Sequence Listing. Such oligonucleotides can be used in conjunction with the polymerase chain reaction (PCR) to screen libraries, isolate clones, and prepare cloning and sequencing templates, etc.

Alternatively, such NHP oligonucleotides can be used as hybridization probes for screening libraries, and assessing gene expression patterns (particularly using a micro array or high-throughput "chip" format). Additionally, a series of the described NHP oligonucleotide sequences, or the complements thereof, can be used to represent all or a portion of the described NHP sequences. The oligonucleotides, typically between about 16 to about 40 (or any whole number within the stated range) nucleotides in length may partially overlap each other and/or a NHP sequence may be represented using oligonucleotides that do not

overlap. Accordingly, the described NHP polynucleotide sequences shall typically comprise at least about two or three distinct oligonucleotide sequences of at least about 18, and preferably about 25, nucleotides in length that are each first disclosed in the described Sequence Listing. Such oligonucleotide sequences may begin at any nucleotide present within a sequence in the Sequence Listing and proceed in either a sense (5'-to-3') orientation vis-a-vis the described sequence or in an antisense orientation.

For oligonucleotide probes, highly stringent conditions may refer, e.g., to washing in 6xSSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos). These nucleic acid molecules may encode or act as NHP gene antisense molecules, useful, for example, in NHP gene regulation (for and/or as antisense primers in amplification reactions of NHP gene nucleic acid sequences). With respect to NHP gene regulation, such techniques can be used to regulate biological functions. Further, such sequences may be used as part of ribozyme and/or triple helix sequences that are also useful for NHP gene regulation.

Inhibitory antisense or double stranded oligonucleotides can additionally comprise at least one modified base moiety which is selected from the group including but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl)uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil,

5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide can also comprise at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

10 In yet another embodiment, the antisense oligonucleotide will comprise at least one modified phosphate backbone selected from the group consisting of a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or
15 analog thereof.

In yet another embodiment, the antisense oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to
20 each other (Gautier *et al.*, 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue *et al.*, 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue *et al.*, 1987, FEBS Lett. 215:327-330).
Alternatively, double stranded RNA can be used to disrupt the
25 expression and function of a targeted NHP.

Oligonucleotides of the invention can be synthesized by standard methods known in the art, *e.g.* by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate
30 oligonucleotides can be synthesized by the method of Stein *et al.* (1988, Nucl. Acids Res. 16:3209), and methylphosphonate oligonucleotides can be prepared by use of controlled pore glass

polymer supports (Sarin *et al.*, 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85:7448-7451), etc.

Low stringency conditions are well known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook *et al.*, 1989, *Molecular Cloning, A Laboratory Manual* (and periodic updates thereof), Cold Springs Harbor Press, N.Y.; and Ausubel *et al.*, 1989, *Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, N.Y.

Alternatively, suitably labeled NHP nucleotide probes can be used to screen a human genomic library using appropriately stringent conditions or by PCR. The identification and characterization of human genomic clones is helpful for identifying polymorphisms (including, but not limited to, nucleotide repeats, microsatellite alleles, single nucleotide polymorphisms, or coding single nucleotide polymorphisms), determining the genomic structure of a given locus/allele, and designing diagnostic tests. For example, sequences derived from regions adjacent to the intron/exon boundaries of the human gene can be used to design primers for use in amplification assays to detect mutations within the exons, introns, splice sites (e.g., splice acceptor and/or donor sites), etc., that can be used in diagnostics and pharmacogenomics.

Further, a NHP homolog can be isolated from nucleic acid from an organism of interest by performing PCR using two degenerate or "wobble" oligonucleotide primer pools designed on the basis of amino acid sequences within the NHP products disclosed herein. The template for the reaction may be total RNA, mRNA, and/or cDNA obtained by reverse transcription of mRNA prepared from human or non-human cell lines or tissue known or suspected to express an allele of a NHP gene. The PCR product can be subcloned and sequenced to ensure that the amplified sequences represent the

sequence of the desired NHP gene. The PCR fragment can then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment can be labeled and used to screen a cDNA library, such as a bacteriophage cDNA library.

- 5 Alternatively, the labeled fragment can be used to isolate genomic clones via the screening of a genomic library.

PCR technology can also be used to isolate full length cDNA sequences. For example, RNA can be isolated, following standard procedures, from an appropriate cellular or tissue source (*i.e.*,

- 10 one known, or suspected, to express a NHP gene, such as, for example, testis tissue). A reverse transcription (RT) reaction can be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid
15 may then be "tailed" using a standard terminal transferase reaction, the hybrid may be digested with RNase H, and second strand synthesis may then be primed with a complementary primer. Thus, cDNA sequences upstream of the amplified fragment can be isolated. For a review of cloning strategies that can be used,
20 see *e.g.*, Sambrook *et al.*, 1989, *supra*.

- A cDNA encoding a mutant NHP gene can be isolated, for example, by using PCR. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known or suspected to be expressed in an
25 individual putatively carrying a mutant NHP allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal gene. Using these two primers, the product is then amplified via PCR,
30 optionally cloned into a suitable vector, and subjected to DNA sequence analysis through methods well known to those of skill in the art. By comparing the DNA sequence of the mutant NHP allele to that of a corresponding normal NHP allele, the mutation(s)

responsible for the loss or alteration of function of the mutant NHP gene product can be ascertained.

Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of or known to carry a mutant NHP allele (e.g., a person manifesting a NHP-associated phenotype such as, for example, obesity, high blood pressure, connective tissue disorders, infertility, etc.), or a cDNA library can be constructed using RNA from a tissue known, or suspected, to express a mutant NHP allele. A normal NHP gene, or any suitable fragment thereof, can then be labeled and used as a probe to identify the corresponding mutant NHP allele in such libraries. Clones containing mutant NHP gene sequences can then be purified and subjected to sequence analysis according to methods well known to those skilled in the art.

Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a tissue known, or suspected, to express a mutant NHP allele in an individual suspected of or known to carry such a mutant allele. In this manner, gene products made by the putatively mutant tissue can be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against normal NHP product, as described below. (For screening techniques, see, for example, Harlow, E. and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor.)

Additionally, screening can be accomplished by screening with labeled NHP fusion proteins, such as, for example, alkaline phosphatase-NHP or NHP-alkaline phosphatase fusion proteins. In cases where a NHP mutation results in an expressed gene product with altered function (e.g., as a result of a missense or a frameshift mutation), polyclonal antibodies to NHP are likely to cross-react with a corresponding mutant NHP gene product. Library clones detected via their reaction with such labeled antibodies

can be purified and subjected to sequence analysis according to methods well known in the art.

The invention also encompasses (a) DNA vectors that contain any of the foregoing NHP coding sequences and/or their complements (i.e., antisense); (b) DNA expression vectors that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences (for example, baculo virus as described in U.S. Patent No. 5,869,336 herein incorporated by reference); (c) genetically engineered host cells that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell; and (d) genetically engineered host cells that express an endogenous NHP gene under the control of an exogenously introduced regulatory element (i.e., gene activation). As used herein, regulatory elements include, but are not limited to, inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate expression. Such regulatory elements include but are not limited to the human cytomegalovirus (hCMV) immediate early gene, regulatable, viral elements (particularly retroviral LTR promoters), the early or late promoters of SV40 adenovirus, the *lac* system, the *trp* system, the *TAC* system, the *TRC* system, the major operator and promoter regions of phage lambda, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase (PGK), the promoters of acid phosphatase, and the promoters of the yeast α -mating factors.

The present invention also encompasses antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists and agonists of a NHP, as well as compounds or nucleotide constructs that inhibit expression of a NHP gene (transcription factor inhibitors, antisense and ribozyme molecules, or gene or regulatory sequence replacement constructs), or promote the expression of a NHP (e.g., expression constructs in which NHP

coding sequences are operatively associated with expression control elements such as promoters, promoter/enhancers, etc.).

The NHPs or NHP peptides, NHP fusion proteins, NHP nucleotide sequences, antibodies, antagonists and agonists can be useful for the detection of mutant NHPs or inappropriately expressed NHPs for the diagnosis of disease. The NHPs or NHP peptides, NHP fusion proteins, NHP nucleotide sequences, host cell expression systems, antibodies, antagonists, agonists and genetically engineered cells and animals can be used for screening for drugs (or high throughput screening of combinatorial libraries) effective in the treatment of the symptomatic or phenotypic manifestations of perturbing the normal function of NHP in the body. The use of engineered host cells and/or animals may offer an advantage in that such systems allow not only for the identification of compounds that bind to the endogenous receptor for a NHP, but can also identify compounds that trigger NHP-mediated activities or pathways.

Finally, the NHP products can be used as therapeutics. For example, soluble derivatives such as NHP peptides/domains corresponding to NHP, NHP fusion protein products (especially NHP-Ig fusion proteins, i.e., fusions of a NHP, or a domain of a NHP, to an IgFc), NHP antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists or agonists (including compounds that modulate or act on downstream targets in a NHP-mediated pathway) can be used to directly treat diseases or disorders. For instance, the administration of an effective amount of soluble NHP, or a NHP-IgFc fusion protein or an anti-idiotypic antibody (or its Fab) that mimics a NHP could activate or effectively antagonize the endogenous NHP receptor. Nucleotide constructs encoding such NHP products can be used to genetically engineer host cells to express such products *in vivo*; these genetically engineered cells function as "bioreactors" in the body delivering a continuous supply of a NHP, a NHP peptide, or a NHP

fusion protein to the body. Nucleotide constructs encoding functional NHP, mutant NHPs, as well as antisense and ribozyme molecules can also be used in "gene therapy" approaches for the modulation of NHP expression. Thus, the invention also encompasses pharmaceutical formulations and methods for treating biological disorders.

Various aspects of the invention are described in greater detail in the subsections below.

5.1 THE NHP SEQUENCES

The cDNA sequences and corresponding deduced amino acid sequences of the described NHPs are presented in the Sequence Listing. SEQ ID NO:7 describes a NHP ORF as well as flanking regions. The NHP nucleotides were obtained from human cDNA libraries using probes and/or primers generated from human gene trapped sequence tags, and genomic sequence. Expression analysis has provided evidence that the described NHP can be expressed a variety of human cells as well as gene trapped human cells.

5.2 NHPs AND NHP POLYPEPTIDES

NHPs, polypeptides, peptide fragments, mutated, truncated, or deleted forms of the NHPs, and/or NHP fusion proteins can be prepared for a variety of uses. These uses include, but are not limited to, the generation of antibodies, as reagents in diagnostic assays, for the identification of other cellular gene products related to a NHP, as reagents in assays for screening for compounds that can be as pharmaceutical reagents useful in the therapeutic treatment of mental, biological, or medical disorders and disease.

The Sequence Listing discloses the amino acid sequences encoded by the described NHP polynucleotides. The NHPs display an initiator methionine in DNA sequence contexts consistent with a translation initiation site, and several of the ORFs display a

consensus signal sequence which can indicate that the described NHP ORFs are secreted proteins, or can be membrane associated.

The NHP amino acid sequences of the invention include the amino acid sequences presented in the Sequence Listing as well as analogues and derivatives thereof. Further, corresponding NHP homologues from other species are encompassed by the invention. In fact, any NHPs encoded by a NHP nucleotide sequence described above are within the scope of the invention, as are any novel polynucleotide sequences encoding all or any novel portion of an amino acid sequence presented in the Sequence Listing. The degenerate nature of the genetic code is well known, and, accordingly, each amino acid presented in the Sequence Listing, is generically representative of the well known nucleic acid "triplet" codon, or in many cases codons, that can encode the amino acid. As such, as contemplated herein, the amino acid sequences presented in the Sequence Listing, when taken together with the genetic code (see, for example, Table 4-1 at page 109 of "Molecular Cell Biology", 1986, J. Darnell et al. eds., Scientific American Books, New York, NY, herein incorporated by reference) are generically representative of all the various permutations and combinations of nucleic acid sequences that can encode such amino acid sequences.

The invention also encompasses proteins that are functionally equivalent to the NHPs encoded by the presently described nucleotide sequences as judged by any of a number of criteria, including, but not limited to, the ability to bind and cleave a substrate of a NHP, or the ability to effect an identical or complementary downstream pathway, or a change in cellular metabolism (e.g., proteolytic activity, ion flux, tyrosine phosphorylation, etc.). Such functionally equivalent NHP proteins include, but are not limited to, additions or substitutions of amino acid residues within the amino acid sequence encoded by the NHP nucleotide sequences described above, but which result in a

silent change, thus producing a functionally equivalent gene product. Amino acid substitutions can be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues

involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

A variety of host-expression vector systems can be used to express the NHP nucleotide sequences of the invention. Where, as in the present instance, a NHP peptide or NHP polypeptide is thought to be a soluble or secreted molecule, the peptide or polypeptide can be recovered from the culture media. Such expression systems also encompass engineered host cells that express NHP, or functional equivalent, *in situ*. Purification or enrichment of a NHP from such expression systems can be accomplished using appropriate detergents and lipid micelles and methods well known to those skilled in the art. However, such engineered host cells themselves may be used in situations where it is important not only to retain the structural and functional characteristics of a NHP, but to assess biological activity, e.g., in drug screening assays.

The expression systems that may be used for purposes of the invention include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing NHP nucleotide sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing NHP encoding nucleotide sequences; insect cell systems infected with recombinant virus expression vectors (e.g.,

baculovirus) containing NHP sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid)

- 5 containing NHP nucleotide sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus
10 7.5K promoter).

- In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the NHP product being expressed. For example, when a large quantity of such a protein is to be produced for the generation of
15 pharmaceutical compositions of and/or containing a NHP, or for raising antibodies to a NHP, vectors that direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J.
20 2:1791), in which a NHP coding sequence may be ligated individually into the vector in frame with the *lacZ* coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke & Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX
25 vectors (Pharmacia or American Type Culture Collection) can also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the
30 presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

In an insect system, *Autographa californica* nuclear polyhidrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. A NHP coding sequence can be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of NHP coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (*i.e.*, virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted sequence is expressed (*e.g.*, see Smith *et al.*, 1983, J. Virol. 46: 584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the NHP nucleotide sequence of interest may be ligated to an adenovirus transcription/translation control complex, *e.g.*, the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (*e.g.*, region E1 or E3) will result in a recombinant virus that is viable and capable of expressing a NHP product in infected hosts (*e.g.*, See Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific initiation signals may also be required for efficient translation of inserted NHP nucleotide sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire NHP sequence or cDNA, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of a NHP coding sequence is inserted, exogenous translational control signals,

including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control
5 signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (See Bittner et al., 1987, Methods in Enzymol. 153:516-544).

10 In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g.,
15 cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct
20 modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in
25 particular, human cell lines.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the NHP sequences described above can be engineered. Rather than using expression vectors which contain
30 viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following

the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows
5 cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express a NHP product. Such engineered cell lines may be particularly useful in screening and evaluation of
10 compounds that affect the endogenous activity of a NHP product.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, Proc.
15 Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy, et al., 1980, Cell 22:817) genes can be employed in tk⁻, hgp⁻ or aprt⁻ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to
20 methotrexate (Wigler, et al., 1980, Natl. Acad. Sci. USA 77:3567; O'Hare, et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al.,
25 1981, J. Mol. Biol. 150:1); and hyg^r, which confers resistance to hygromycin (Santerre, et al., 1984, Gene 30:147).

Alternatively, any fusion protein can be readily purified by utilizing an antibody specific for the fusion protein being expressed. For example, a system described by Janknecht et al.
30 allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, Proc. Natl. Acad. Sci. USA 88:8972-8976). In this system, the gene of

interest is subcloned into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded
5 onto Ni²⁺-nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

5.3 ANTIBODIES TO NHP PRODUCTS

Antibodies that specifically recognize one or more epitopes
10 of a NHP, or epitopes of conserved variants of a NHP, or peptide fragments of a NHP are also encompassed by the invention. Such antibodies include but are not limited to polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments,
15 fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

The antibodies of the invention may be used, for example, in the detection of a NHP in a biological sample and may, therefore,
20 be utilized as part of a diagnostic or prognostic technique whereby patients may be tested for abnormal amounts of NHP. Such antibodies may also be utilized in conjunction with, for example, compound screening schemes for the evaluation of the effect of test compounds on expression and/or activity of a NHP sequence
25 product. Additionally, such antibodies can be used in conjunction gene therapy to, for example, evaluate the normal and/or engineered NHP-expressing cells prior to their introduction into the patient. Such antibodies may additionally be used as a method for the inhibition of abnormal NHP activity. Thus, such
30 antibodies may, therefore, be utilized as part of treatment methods.

For the production of antibodies, various host animals may be immunized by injection with a NHP, an NHP peptide (e.g., one

corresponding to a functional domain of a NHP), truncated NHP polypeptides (NHP in which one or more domains have been deleted), functional equivalents of a NHP or mutated variants of a NHP. Such host animals may include but are not limited to pigs, rabbits, mice, goats, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's adjuvant (complete and incomplete), mineral salts such as aluminum hydroxide or aluminum phosphate, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*. Alternatively, the immune response could be enhanced by combination and or coupling with molecules such as keyhole limpet hemocyanin, tetanus toxoid, diptheria toxoid, ovalbumin, cholera toxin or fragments thereof. Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of the immunized animals.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, can be obtained by any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique of Kohler and Milstein, (1975, Nature 256:495-497; and U.S. Patent No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. USA 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, Monoclonal Antibodies And Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated in vitro or in vivo. Production of high titers of mAbs in vivo makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison *et al.*, 1984, Proc. Natl. Acad. Sci., 81:6851-6855; Neuberger *et al.*, 1984, Nature, 312:604-608; Takeda *et al.*, 1985, Nature, 314:452-454) by splicing the genes
5 from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived
10 from a murine mAb and a human immunoglobulin constant region. Such technologies are described in U.S. Patents Nos. 6,075,181 and 5,877,397 and their respective disclosures which are herein incorporated by reference in their entirety.

Alternatively, techniques described for the production of
15 single chain antibodies (U.S. Patent 4,946,778; Bird, 1988, Science 242:423-426; Huston *et al.*, 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; and Ward *et al.*, 1989, Nature 334:544-546) can be adapted to produce single chain antibodies against NHP sequence products. Single chain antibodies are formed by linking the heavy
20 and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, such fragments include, but are not limited to: the F(ab')₂ fragments which can
25 be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed (Huse *et al.*, 1989, Science, 246:1275-1281) to allow rapid and easy identification of
30 monoclonal Fab fragments with the desired specificity.

Antibodies to a NHP can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" a given NHP, using techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, 1993, FASEB J 7(5):437-444; and Nissinoff, 1991, J. Immunol. 147(8):2429-2438). For example antibodies which bind to a NHP domain and competitively inhibit the binding of NHP to its cognate receptor can be used to generate anti-idiotypes that "mimic" a NHP and, therefore, bind and activate or neutralize a receptor. Such anti-idiotypic antibodies or Fab fragments of such anti-idiotypes can be used in therapeutic regimens involving a NHP signaling pathway.

The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims. All cited publications, patents, and patent applications are herein incorporated by reference in their entirety.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising at least 24 contiguous bases of nucleotide sequence first disclosed
5 in the NHP sequence described in SEQ ID NO: 3.

2. An isolated nucleic acid molecule comprising a nucleotide sequence that:

- 10 (a) encodes the amino acid sequence shown in SEQ ID NO: 4; and
(b) hybridizes under stringent conditions to the nucleotide sequence of SEQ ID NO: 3 or the complement thereof.

15 3. An isolated nucleic acid molecule according to Claim 1 wherein said nucleotide sequence is present in cDNA.

4. An isolated nucleic acid molecule encoding the amino acid sequence presented in SEQ ID NO:2.
20

5. An isolated nucleic acid molecule encoding the amino acid sequence presented in SEQ ID NO:6.

ABSTRACT

Novel human polynucleotide and polypeptide sequences are disclosed that can be used in therapeutic, diagnostic, and pharmacogenomic applications.

PATENT APPLICATION

DECLARATION AND POWER OF ATTORNEY FOR PATENT APPLICATION

ATTORNEY DOCKET NO. LEX-0071-USA

As a below named inventor, I hereby declare that:

My residence/post office address and citizenship are as stated below next to my name;

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Novel Human Proteins and Polynucleotides Encoding the Same

the specification of which is attached hereto unless the following box is checked:

() was filed on _____ as US Application Serial No. or PCT International Application

Number _____ and was amended on _____ (if applicable).

I hereby state that I have reviewed and understood the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above. I acknowledge the duty to disclose all information which is material to patentability as defined in 37 CFR 1.56.

Foreign Application(s) and/or Claim of Foreign Priority

I hereby claim foreign priority benefits under Title 35, United States Code Section 119 of any foreign application(s) for patent or inventor(s) certificate listed below and have also identified below any foreign application for patent or inventor(s) certificate having a filing date before that of the application on which priority is claimed:

COUNTRY	APPLICATION NUMBER	DATE FILED	PRIORITY CLAIMED UNDER 35 U.S.C. 119
			YES: _____ NO: _____
			YES: _____ NO: _____

Provisional Application

I hereby claim the benefit under Title 35, United States Code Section 119(e) of any United States provisional application(s) listed below:

APPLICATION SERIAL NUMBER	FILING DATE
60/160,285	10/19/1999
60/183,583	02/18/2000

U.S. Priority Claim

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

APPLICATION SERIAL NUMBER	FILING DATE	STATUS(patented/pending/abandoned)

POWER OF ATTORNEY:

As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) listed below to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Lance K. Ishimoto, Reg. No. 41866

Send Correspondence to:

Lance K. Ishimoto
Lexicon Genetics Incorporated
4000 Research Forest Drive
The Woodlands, TX 77381

Direct Telephone Calls To:

Lance K. Ishimoto
(281) 362-6554

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION (continued)

ATTORNEY DOCKET NO. LEX-0071-USA

Full Name of Inventor: Gregory Donoho Citizenship: USA

Residence: 41 Fallenstone Dr., The Woodlands, TX 77381

Post Office Address: Same

Inventor's Signature _____ Date _____

Full Name of Inventor: C. Alexander Turner, Jr. Citizenship: USA

Residence: 67 Winter Wheat Place, The Woodlands, TX 77381

Post Office Address: Same

Inventor's Signature _____ Date _____

Full Name of Inventor: Michael C. Nehls Citizenship: Germany

Residence: Paul-Keller-Strasse 6, Stockdorf, Germany 82131

Post Office Address: Same

Inventor's Signature _____ Date _____

Full Name of Inventor: Glenn Friedrich Citizenship: Canada

Residence: c/o Breland & Breland, Houston, TX 77004

Post Office Address: Same

Inventor's Signature _____ Date _____

Full Name of Inventor: Brian Zambrowicz Citizenship: USA

Residence: 18 Firethorne Place, The Woodlands, TX 77382

Post Office Address: Same

Inventor's Signature _____ Date _____

Full Name of Inventor: Arthur T. Sands Citizenship: USA

Residence: 163 Bristol Bend Circle, The Woodlands, TX 77382

Post Office Address: Same

Inventor's Signature _____ Date _____

SEQUENCE LISTING

<110> Donoho, Gregory
Turner, C. Alexander Jr.
Nehls, Michael
Friedrich, Glenn
Zambrowicz, Brian
Sands, Arthur T.

<120> Novel Human Proteins and Polynucleotides
Encoding the Same

<130> LEX-0071-USA

<150> US 60/160,285

<151> 1999-10-19

<150> US 60/183,583

<151> 2000-02-18

<160> 7

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1464

<212> DNA

<213> homo sapiens

<400> 1

atgacatcta	agaattatcc	cgggacctac	cccaatcaca	ctgtttgcga	aaagacaatt	60
acagtaccaa	aggggaaaag	actgattctg	aggtttgggag	atattggatat	cgaatcccag	120
acctgtgctt	ctgactatct	tctcttcacc	agctcttcag	atcaatatgg	tccatactgt	180
ggaagtatga	ctgttcccaa	agaactcttg	ttgaacacaa	gtgaagtaac	cgctcgcttt	240
gagagtgat	cccacatttc	tggccggggt	tttttgcgtga	cctatgcgag	cagcgaccat	300
ccagatttaa	taacatgttt	ggaacgagct	agccattatt	tgaagacaga	atacagcaaa	360
ttctgccccg	ctggttgtag	agacgtagca	ggagacattt	ctgggaatat	ggtagatgga	420
tatagagata	cctctttatt	gtgcaaagct	gccatccatg	caggaataat	tgctgatgaa	480
ctaggtggcc	agatcagtgt	gcttcagcgc	aaagggatca	gtcgatatga	agggattctg	540
gccaatgggt	ttcttttcgag	ggatgggtcc	ctgtcagaca	agcgatttct	gtttacctcc	600
aatgggttga	gcagatccct	gagttttgaa	cctgacgggc	aaatcagagc	ttcttctctca	660
tggcagtcg	tcaatgagag	tggagaccaa	gttcactggt	ctctgggcca	agcccgactt	720
caggaccaag	gcccatcatg	ggcttcgggc	gacagtagca	acaaccacaa	accacgagag	780
tggctggaga	tcgatttggt	ggagaaaaag	aaaataacag	gaattaggac	cacaggatct	840
acacagtcga	acttcaactt	ttatgttaag	agttttgtga	tgaacttcaa	aaaacaataat	900
tctaagtgtga	agacctataa	aggaattgtg	aataatgaag	aaaagggtgt	tcagggttaac	960
tctaactttc	gggacccagt	gcaaaaacaat	ttcatccctc	ccatcgtggc	cagatatgtg	1020
cgggtgtgct	ccagagactt	gcaccagagg	atagccttga	aggtggagct	cattggttgc	1080
cagattacac	aaggtaatga	ttcatgtggt	tggcgcaaga	caagtcaaaag	caccagtggt	1140
tcaactaaga	aagaagatga	gacaatcaca	aggcccattc	cctcggaaga	aacatccaca	1200
ggataaaaac	ttacaacggt	ggctattcca	ttggtgctcc	tgtgtgtctt	ggattttgct	1260
ggattgggga	tctttgcagc	ctttagaaga	aagaagaaga	aaggaagtc	gtatggatca	1320
cgggaggctc	agaaaacaga	ctgttgggaag	cagattaaat	atccctttgc	cagacatcag	1380
tcagctgagt	ttaccatcag	ctatgataat	gagaaggaga	tgacacaaaa	gttagatctc	1440
atcacaaagt	atatggcagg	ttaa				1464

<210> 2
 <211> 487
 <212> PRT
 <213> homo sapiens

<400> 2
 Met Thr Ser Lys Asn Tyr Pro Gly Thr Tyr Pro Asn His Thr Val Cys
 1 5 10 15
 Glu Lys Thr Ile Thr Val Pro Lys Gly Lys Arg Leu Ile Leu Arg Leu
 20 25 30
 Gly Asp Leu Asp Ile Glu Ser Gln Thr Cys Ala Ser Asp Tyr Leu Leu
 35 40 45
 Phe Thr Ser Ser Ser Asp Gln Tyr Gly Pro Tyr Cys Gly Ser Met Thr
 50 55 60
 Val Pro Lys Glu Leu Leu Asn Thr Ser Glu Val Thr Val Arg Phe
 65 70 75 80
 Glu Ser Gly Ser His Ile Ser Gly Arg Gly Phe Leu Leu Thr Tyr Ala
 85 90 95
 Ser Ser Asp His Pro Asp Leu Ile Thr Cys Leu Glu Arg Ala Ser His
 100 105 110
 Tyr Leu Lys Thr Glu Tyr Ser Lys Phe Cys Pro Ala Gly Cys Arg Asp
 115 120 125
 Val Ala Gly Asp Ile Ser Gly Asn Met Val Asp Gly Tyr Arg Asp Thr
 130 135 140
 Ser Leu Leu Cys Lys Ala Ala Ile His Ala Gly Ile Ile Ala Asp Glu
 145 150 155 160
 Leu Gly Gly Gln Ile Ser Val Leu Gln Arg Lys Gly Ile Ser Arg Tyr
 165 170 175
 Glu Gly Ile Leu Ala Asn Gly Val Leu Ser Arg Asp Gly Ser Leu Ser
 180 185 190
 Asp Lys Arg Phe Leu Phe Thr Ser Asn Gly Cys Ser Arg Ser Leu Ser
 195 200 205
 Phe Glu Pro Asp Gly Gln Ile Arg Ala Ser Ser Ser Trp Gln Ser Val
 210 215 220
 Asn Glu Ser Gly Asp Gln Val His Trp Ser Pro Gly Gln Ala Arg Leu
 225 230 235 240
 Gln Asp Gln Gly Pro Ser Trp Ala Ser Gly Asp Ser Ser Asn Asn His
 245 250 255
 Lys Pro Arg Glu Trp Leu Glu Ile Asp Leu Gly Glu Lys Lys Ile
 260 265 270
 Thr Gly Ile Arg Thr Thr Gly Ser Thr Gln Ser Asn Phe Asn Phe Tyr
 275 280 285
 Val Lys Ser Phe Val Met Asn Phe Lys Asn Asn Asn Ser Lys Trp Lys
 290 295 300
 Thr Tyr Lys Gly Ile Val Asn Asn Glu Glu Lys Val Phe Gln Gly Asn
 305 310 315 320
 Ser Asn Phe Arg Asp Pro Val Gln Asn Asn Phe Ile Pro Pro Ile Val
 325 330 335
 Ala Arg Tyr Val Arg Val Val Pro Gln Thr Trp His Gln Arg Ile Ala
 340 345 350
 Leu Lys Val Glu Leu Ile Gly Cys Gln Ile Thr Gln Gly Asn Asp Ser
 355 360 365
 Leu Val Trp Arg Lys Thr Ser Gln Ser Thr Ser Val Ser Thr Lys Lys
 370 375 380
 Glu Asp Glu Thr Ile Thr Arg Pro Ile Pro Ser Glu Glu Thr Ser Thr
 385 390 395 400

Gly Ile Asn Ile Thr Thr Val Ala Ile Pro Leu Val Leu Leu Val Val
 405 410 415
 Leu Val Phe Ala Gly Met Gly Ile Phe Ala Ala Phe Arg Lys Lys Lys
 420 425 430
 Lys Lys Gly Ser Pro Tyr Gly Ser Ala Glu Ala Gln Lys Thr Asp Cys
 435 440 445
 Trp Lys Gln Ile Lys Tyr Pro Phe Ala Arg His Gln Ser Ala Glu Phe
 450 455 460
 Thr Ile Ser Tyr Asp Asn Glu Lys Glu Met Thr Gln Lys Leu Asp Leu
 465 470 475 480
 Ile Thr Ser Asp Met Ala Gly
 485

<210> 3
 <211> 1761
 <212> DNA
 <213> homo sapiens

<400> 3
 atgggattcg gtgcggggca ggcactgcgc cccgtccggc gcgcgcgctc gtccgcagag 60
 gagggcgccc ggcccgggca gctgcggcto gggatccgct gaggggaggg cgagcttgcc 120
 aagctggggc ccagcgggggt catggtgccc ggcgcccgcg gcggcgcgcg actggcgcg 240
 gctgcggggc ggggcctcct ggctttgtcg ctgcgggtct cgcgcccgct ccggctgcag 360
 gcggaggagc tgggtgatgg ctgtggacac ctagtgaact atcaggatag tggcacaatg 300
 acatctaaaga attatccggg gacctacccc aatcacactg ttgcgaaaaa gacaattaca 360
 gtaccaaaagg gaaaagaagt gattctgagg ttgggagatt tggatatgca atccagagc 420
 tgtgtctctg acctatctct ctccaccagc tcttcagatc aatatgggtc atactgtgga 480
 agtatgactg ttccc aaaaga actcttgggt aacacaagtg aagtaaccgt ccgctttgag 540
 agtggatccc acatttctgg ccgggggttt ttgctgacct atgcgagcag cgaccatcca 600
 gatttaataa catgttttga acgagctagc cattatttga agacagaata cagcaaatcc 660
 tgcccagctg gttgtagaga cgtagcagga gacatttctg ggaatatggt agatggatat 720
 agagatacct ctttattgtg caaagctgcc atccatgcag gaataattgc tgatgaacta 780
 ggtggccaga tcagtgtgct tcagcgcaaa gggatcagtc gatatgaagg gattctggcc 840
 aatggtgttc ttctgaggga tgggtccctg tcagacaagc gatttctgtt tacctccaat 900
 ggttgacgca gacccctgag ttttgaacct gacgggcaaa tcagagcttc ttccctcatg 960
 cagtcgggtca atgagagtgg agaccaagtt cactgtgtct ctggccaagc ccgacttcag 1020
 gaccaaggcc catcatgggc ttgcggcgac agtagcaaca accacaacc acgagagtgg 1080
 ctgtagatgc atttggggga gaaaaagaaa ataacaggaa tttagaccac aggatctaca 1140
 cagtcgaact tcaactttta tggtaaagat tttgtgatga acttcaaaaa caataattct 1200
 aagtggaaaga cctataaaagg aattgtgaa aatgaagaaa aggtgtttca gggtaactct 1260
 aactttcggg acccagtgca aaacaatttc atccctccca tegtggccag atatgtgagg 1320
 gttgtccccc agacatggca ccagaggata gccttgaagg tggagctcat tggttgccag 1380
 attacacaag gtaatgattc attggtgtgg cgcaagacaa gtcaagcac cagtggttca 1440
 actaagaaaag aagatgagac aatcacaaag cccatccctc cggaagaaac atccacagga 1500
 ataaaacatta caacggtggc tattccattg gtgctccttg ttgtcctggt gtttctggtg 1560
 atgggggatct ttgcagcctt tagaaaagag aagaagaaa gaagtcocga tggatcagcg 1620
 gaggtccaga aaacagactg ttggaagcag attaaatcat cctttgcgag acatcagtca 1680
 gctgagttaa ccatcagcta tgataatgag aaggagatga cacaaaaagt agatctcatc 1740
 acaagtgaata tggcaggtta a 1761

<210> 4
 <211> 586
 <212> PRT
 <213> homo sapiens

<400> 4

Met Gly Phe Gly Ala Gly Gln Arg Leu Arg Pro Val Pro Ala Pro Arg
 1 5 10 15
 Ser Ser Ala Glu Ala Ala Arg Pro Gly Gln Leu Arg Leu Gly Ile
 20 25 30
 Arg Arg Gly Glu Ala Glu Leu Ala Lys Leu Ala Pro Ser Gly Val Met
 35 40 45
 Val Pro Gly Ala Arg Gly Gly Ala Leu Ala Arg Ala Ala Gly Arg
 50 55 60
 Gly Leu Leu Ala Leu Leu Leu Ala Val Ser Ala Pro Leu Arg Leu Gln
 65 70 75 80
 Ala Glu Glu Leu Gly Asp Gly Cys Gly His Leu Val Thr Tyr Gln Asp
 85 90 95
 Ser Gly Thr Met Thr Ser Lys Asn Tyr Pro Gly Thr Tyr Pro Asn His
 100 105 110
 Thr Val Cys Glu Lys Thr Ile Thr Val Pro Lys Gly Lys Arg Leu Ile
 115 120 125
 Leu Arg Leu Gly Asp Leu Asp Ile Glu Ser Gln Thr Cys Ala Ser Asp
 130 135 140
 Tyr Leu Leu Phe Thr Ser Ser Ser Asp Gln Tyr Gly Pro Tyr Cys Gly
 145 150 155 160
 Ser Met Thr Val Pro Lys Glu Leu Leu Leu Asn Thr Ser Glu Val Thr
 165 170 175
 Val Arg Phe Glu Ser Gly Ser His Ile Ser Gly Arg Gly Phe Leu Leu
 180 185 190
 Thr Tyr Ala Ser Ser Asp His Pro Asp Leu Ile Thr Cys Leu Glu Arg
 195 200 205
 Ala Ser His Tyr Leu Lys Thr Glu Tyr Ser Lys Phe Cys Pro Ala Gly
 210 215 220
 Cys Arg Asp Val Ala Gly Asp Ile Ser Gly Asn Met Val Asp Gly Tyr
 225 230 235 240
 Arg Asp Thr Ser Leu Leu Cys Lys Ala Ala Ile His Ala Gly Ile Ile
 245 250 255
 Ala Asp Glu Leu Gly Gly Gln Ile Ser Val Leu Gln Arg Lys Gly Ile
 260 265 270
 Ser Arg Tyr Glu Gly Ile Leu Ala Asn Gly Val Leu Ser Arg Asp Gly
 275 280 285
 Ser Leu Ser Asp Lys Arg Phe Leu Phe Thr Ser Asn Gly Cys Ser Arg
 290 295 300
 Ser Leu Ser Phe Glu Pro Asp Gly Gln Ile Arg Ala Ser Ser Ser Trp
 305 310 315 320
 Gln Ser Val Asn Glu Ser Gly Asp Gln Val His Trp Ser Pro Gly Gln
 325 330 335
 Ala Arg Leu Gln Asp Gln Gly Pro Ser Trp Ala Ser Gly Asp Ser Ser
 340 345 350
 Asn Asn His Lys Pro Arg Glu Trp Leu Glu Ile Asp Leu Gly Glu Lys
 355 360 365
 Lys Lys Ile Thr Gly Ile Arg Thr Thr Gly Ser Thr Gln Ser Asn Phe
 370 375 380
 Asn Phe Tyr Val Lys Ser Phe Val Met Asn Phe Lys Asn Asn Asn Ser
 385 390 395 400
 Lys Trp Lys Thr Tyr Lys Gly Ile Val Asn Asn Glu Glu Lys Val Phe
 405 410 415
 Gln Gly Asn Ser Asn Phe Arg Asp Pro Val Gln Asn Asn Phe Ile Pro
 420 425 430
 Pro Ile Val Ala Arg Tyr Val Arg Val Val Pro Gln Thr Trp His Gln
 435 440 445

Arg Ile Ala Leu Lys Val Glu Leu Ile Gly Cys Gln Ile Thr Gln Gly
 450 455 460
 Asn Asp Ser Leu Val Trp Arg Lys Thr Ser Gln Ser Thr Ser Val Ser
 465 470 475 480
 Thr Lys Lys Glu Asp Glu Thr Ile Thr Arg Pro Ile Pro Ser Glu Glu
 485 490 495
 Thr Ser Thr Gly Ile Asn Ile Thr Thr Val Ala Ile Pro Leu Val Leu
 500 505 510
 Leu Val Val Leu Val Phe Ala Gly Met Gly Ile Phe Ala Ala Phe Arg
 515 520 525
 Lys Lys Lys Lys Lys Gly Ser Pro Tyr Gly Ser Ala Glu Ala Gln Lys
 530 535 540
 Thr Asp Cys Trp Lys Gln Ile Lys Tyr Pro Phe Ala Arg His Gln Ser
 545 550 555 560
 Ala Glu Phe Thr Ile Ser Tyr Asp Asn Glu Lys Glu Met Thr Gln Lys
 565 570 575
 Leu Asp Leu Ile Thr Ser Asp Met Ala Gly
 580 585

<210> 5

<211> 1620

<212> DNA

<213> homo sapiens

<400> 5

atgggtccccg	gcgcccgcgg	cgccggcgca	ctggcgcggg	ctgcggggcg	gggcctcctg	60
gctttgtctgc	tcgcgggtctc	cgccccgcctc	cggtgcagg	cgaggagagct	gggtgatggc	120
tgtggacacc	tagtgactta	tcaggatagt	ggcacaaatga	catctaaaga	tatccccggg	180
acctacccca	atcacactgt	tgcgaaaaag	acaattacag	tacaaaaggg	gaaaagactg	240
attctcaggt	tggggagattt	ggatatcgaa	tcccagacct	gtgctcttcga	ctatctcttc	300
ttccaccagct	cttcagatca	atatgggtcca	tactgtggaa	gtatgactgt	tcccaaagaa	360
ctcttggtga	acacaagtga	agtaaccgtc	cgctttgaga	gtggatccca	catttctggc	420
cggggttttt	tgtctgacct	tgcgagcagc	gaccatccag	atttaataac	atgtttggaa	480
cgagctagcc	attatttgaa	gacagaatac	agcaaatctt	gcccagctgg	ttgtagagac	540
gtagcaggag	acatttctgg	gaatatggta	gatggatata	gagataacct	tttatgttgc	600
aaagctgcca	tcctatgcag	aataattgct	gatgaactag	gtggccagat	cagtgtgctt	660
cagcgcaaa	ggatcagtcg	atatgaaggg	attctggcca	atgggtgtct	ttcgagggat	720
gggtccctgt	cagacaagcg	atctctgttt	acctccaatg	gttcgagcag	atccttgagt	780
tttgaacctg	acgggcaaat	cagagcttct	tcctcatggc	agtcgggtcaa	tgagaagtga	840
gaccaagttc	actgtgtctc	tggccaagcc	cgacttcagg	accaaggccc	atcatgggct	900
tggggcgaca	gtagcaacaa	ccacaaacca	cgagagtggc	tggagatoga	tttggggggg	960
aaaaaagaaa	taacagggaat	taggaaccaca	ggatctacac	agtcgaactt	caacttttat	1020
gttaagagtt	ttgtgatgaa	cttcaaaaac	aataattcta	agtggaaagac	ctataaaagga	1080
attgtgaata	atgaagaaaa	gggtgtttcag	ggtaactcta	actttcgga	ccagtgcaa	1140
aacaatttca	tccttcccat	cgtggccaga	tatgtgcggg	ttgtccccca	gacatggcac	1200
cagaggatag	ccctgaaggt	ggagctcatt	ggttgccaga	ttacacaagg	taatgattca	1260
ttggtgtggc	gcaagacaag	tcaaagcacc	agtggtttcaa	ctaagaaaga	agatgagaca	1320
atcacaggc	ccatccccctc	ggaagaaaca	tccacaggaa	taaacattac	aacggtggct	1380
attccattgg	tgtcctctgt	tgtcctgggt	ttgtctggaa	tggggacttt	cacagccttt	1440
agaaagaaga	agaagaagg	aagtccttat	ggatcagcgg	aggctcagaa	aacagactgt	1500
tggaagcaga	ttaaattatc	ctttgccaga	catcagtcag	ctcagtttac	catcagctat	1560
gataatgaga	aggagatgac	acaaaagtta	gatctcatca	caagtgatata	ggcaggttaa	1620

<210> 6

<211> 539

<212> PRT

<213> homo sapiens

<400> 6

```

Met Val Pro Gly Ala Arg Gly Gly Gly Ala Leu Ala Arg Ala Ala Gly
1      5      10      15
Arg Gly Leu Leu Ala Leu Leu Ala Val Ser Ala Pro Leu Arg Leu
20      25      30
Gln Ala Glu Glu Leu Gly Asp Gly Cys Gly His Leu Val Thr Tyr Gln
35      40      45
Asp Ser Gly Thr Met Thr Ser Lys Asn Tyr Pro Gly Thr Tyr Pro Asn
50      55      60
His Thr Val Cys Glu Lys Thr Ile Thr Val Pro Lys Gly Lys Arg Leu
65      70      75      80
Ile Leu Arg Leu Gly Asp Leu Asp Ile Glu Ser Gln Thr Cys Ala Ser
85      90      95
Asp Tyr Leu Leu Phe Thr Ser Ser Ser Asp Gln Tyr Gly Pro Tyr Cys
100      105      110
Gly Ser Met Thr Val Pro Lys Glu Leu Leu Asn Thr Ser Glu Val
115      120      125
Thr Val Arg Phe Glu Ser Gly Ser His Ile Ser Gly Arg Gly Phe Leu
130      135      140
Leu Thr Tyr Ala Ser Ser Asp His Pro Asp Leu Ile Thr Cys Leu Glu
145      150      155      160
Arg Ala Ser His Tyr Leu Lys Thr Glu Tyr Ser Lys Phe Cys Pro Ala
165      170      175
Gly Cys Arg Asp Val Ala Gly Asp Ile Ser Gly Asn Met Val Asp Gly
180      185      190
Tyr Arg Asp Thr Ser Leu Leu Cys Lys Ala Ala Ile His Ala Gly Ile
195      200      205
Ile Ala Asp Glu Leu Gly Gly Gln Ile Ser Val Leu Gln Arg Lys Gly
210      215      220
Ile Ser Arg Tyr Glu Gly Ile Leu Ala Asn Gly Val Leu Ser Arg Asp
225      230      235      240
Gly Ser Leu Ser Asp Lys Arg Phe Leu Phe Thr Ser Asn Gly Cys Ser
245      250      255
Arg Ser Leu Ser Phe Glu Pro Asp Gly Gln Ile Arg Ala Ser Ser Ser
260      265      270
Trp Gln Ser Val Asn Glu Ser Gly Asp Gln Val His Trp Ser Pro Gly
275      280      285
Gln Ala Arg Leu Gln Asp Gln Gly Pro Ser Trp Ala Ser Gly Asp Ser
290      295      300
Ser Asn Asn His Lys Pro Arg Glu Trp Leu Glu Ile Asp Leu Gly Glu
305      310      315      320
Lys Lys Lys Ile Thr Gly Ile Arg Thr Thr Gly Ser Thr Gln Ser Asn
325      330      335
Phe Asn Phe Tyr Val Lys Ser Phe Val Met Asn Phe Lys Asn Asn Asn
340      345      350
Ser Lys Trp Lys Thr Tyr Lys Gly Ile Val Asn Asn Gln Glu Lys Val
355      360      365
Phe Gln Gly Asn Ser Asn Phe Arg Asp Pro Val Gln Asn Asn Phe Ile
370      375      380
Pro Pro Ile Val Ala Arg Tyr Val Arg Val Val Pro Gln Thr Trp His
385      390      395      400
Gln Arg Ile Ala Leu Lys Val Glu Leu Ile Gly Cys Gln Ile Thr Gln
405      410      415
Gly Asn Asp Ser Leu Val Trp Arg Lys Thr Ser Gln Ser Thr Ser Val

```

420 425 430
 Ser Thr Lys Lys Glu Asp Glu Thr Ile Thr Arg Pro Ile Pro Ser Glu
 435 440 445
 Glu Thr Ser Thr Gly Ile Asn Ile Thr Thr Val Ala Ile Pro Leu Val
 450 455 460
 Leu Leu Val Val Leu Val Phe Ala Gly Met Gly Ile Phe Ala Ala Phe
 465 470 475 480
 Arg Lys Lys Lys Lys Gly Ser Pro Tyr Gly Ser Ala Glu Ala Gln
 485 490 495
 Lys Thr Asp Cys Trp Lys Gln Ile Lys Tyr Pro Phe Ala Arg His Gln
 500 505 510
 Ser Ala Glu Phe Thr Ile Ser Tyr Asp Asn Glu Lys Glu Met Thr Gln
 515 520 525
 Lys Leu Asp Leu Ile Thr Ser Asp Met Ala Gly
 530 535

<210> 7

<211> 1768

<212> DNA

<213> homo sapiens

<400> 7

ggcggaggag	ctgggtgatg	gctgtggaca	cctagtgtact	tatcaggata	gtggcacaaat	60
gacatctaag	aattatcccg	ggacctacc	caatcacact	gtttgcgaaa	agacaattac	120
agtacaaaag	gggaaaagac	tgattctgag	gttgggagat	ttggatatcg	aatcccgagc	180
ctgtgctttc	gactatcttc	tcttcaccag	ctcttcagat	caatatgtgc	catactgtgg	240
aagtatgact	gttcccaaag	aactcttgtt	gaacacaagt	gaagttaacc	tccgttttga	300
gagtgatgcc	cacattttctg	gccgggggtt	tttgctgacc	tatgcgagca	gcgaccatcc	360
agatttaata	acatgttttg	aacgagctag	ccattatttg	aagacagaat	acagcaaaat	420
ctgcccagct	ggttgttagag	acgtagcagg	agacattttc	gggaatatgg	tagatggata	480
tagagatacc	tctttattgt	gcaaagctgc	catccatgca	ggaataattg	ctgatgaact	540
agggtgccag	atcagltgtc	ttcagcgcga	agggatcagt	cgatatgaag	ggattctggc	600
caatgtgtgt	ctttcgaggg	atgggttccct	gtcagacaa	cgatttctgt	ttacctccaa	660
tggttcgagc	agatccttga	gttttgaacc	tgacggggcaa	atcagagctt	cttctctcat	720
gcagtcggtc	aatgagatgt	gagaccaagt	tcactgggtc	cctggccaa	cccgaactta	780
ggaccaagcg	ccatcatggg	cttcggggca	cagtagcaac	aaccacaaac	cacgagagtg	840
gctggagatc	gatttggggg	agaaaaagaa	aataacagga	attaggacca	cagatctac	900
acagtcgaac	ttcaactttt	atgttaagag	ttttgtgatg	aacttcaaaa	acaataattc	960
taagtggag	acctataaag	gaattgtgaa	taatgaagaa	aaggtgtttc	aggttaactc	1020
taactttcgg	gaccacagtc	aaaaacaatt	catccctccc	atcgtggcca	gatatgtgcg	1080
ggttgtcccc	cagacatggc	accagaggat	agccttgaag	gtggagctca	ttggttgcca	1140
gattacacaa	ggtaaatgatt	catlgtgtgt	gcgcaagaca	agtcacaaagc	ccagtgtttc	1200
aactaagaaa	gaagatgaga	caatcacaag	gcccatcccc	tcggaagaaa	catccacagg	1260
aataaacatt	acaacggtgg	ctatttcatt	gggtctcctt	gttgctcctt	tggttctgtg	1320
aatggggatc	tttgcagcct	ttagaagaaa	gaagaagaaa	ggaagtccgt	atggatcagc	1380
ggaggtctag	aaaaacagat	gttggaaagca	gattaaatat	ccctttgcca	gacatcagtc	1440
agctgcagtt	acacatcagct	atgataatga	gaagggagatg	acacaaaagt	tagatctcat	1500
cacagatgat	atggcaggtt	aactccgttg	actgccaaaa	tagcatcccc	aacgtgcagc	1560
cctccgcttc	tatccagcag	ttgcccggga	tggatctcag	agatgatgat	tggaacacca	1620
tgtttctttc	caccctaaca	acaacaaagg	gcagtaaat	aaagtactct	ttgtaaggta	1680
cagttaccga	ttaactctaga	gataaaatat	tttcttaaaa	atatatttca	ttaaacacct	1740
atgctgtctc	tatgcacaaa	aaaaaaaa				1768